

HIV & HEPATITIS C COINFECTION

DATA REPORTED THROUGH JUNE 30, 2021

New HIV and HCV Diagnoses in Maryland

During 2016-2020, 76 people were newly diagnosed with both HIV and HCV within a 12-month period.

HIV/HCV Diagnoses, 2016-2020	No.	% of Total	Rate	HIV/HCV Diagnoses by Year						
Total	76	100.0%	1.5	30						
Late HIV Diagnosis	21	27.6%		x 25						
Linked to HIV Care in 1 mo.	62	81.6%								
Linked to HIV Care in 3 mo.	70	92.1%		2 0 2.4%						
Age at Diagnosis				25 20 15 1.7% 1.6% 1.5% 0.3%						
13-24	2	2.6%	0.2	1.7%						
25-34	19	25.0%	2.3	₹ 10						
35-44	17	22.4%	2.1	〒 5 0.3%						
45-54	19	25.0%	2.4							
55+	19	25.0%	1.1	2016 2017 2018 2019 2020						
Sex at Birth				Year of HIV Diagnosis						
Male	53	69.7%	2.2							
Female	23	30.3%	0.9	Number of HCV/HIV Diagnoses						
Race/Ethnicity				Percent of Total HIV Diagnoses						
Hispanic	4	5.3%	0.8							
Non-Hispanic Black	53	69.7%	3.5							
Non-Hispanic White	16	21.1%	0.6	Diagnoses by Age at Diagnosis, Sex at						
Non-Hispanic Other	3	3.9%	0.6	Birth and HIV Exposure Category						
Country of Birth				ğ 16 ————						
United States	72	94.7%	1.7	So 14						
Foreign Born	4	5.3%	0.5	<u>12</u> <u>12</u> <u>12</u> <u>12</u> <u>12</u> <u>12</u> <u>12</u> <u>12</u>						
Reported HIV Exposure Catego	ry			Number of Diagnoses						
MSM	16	21.1%		a 8						
IDU	25	32.9%								
MSM/IDU	7	9.2%								
HET	17	22.4%								
Other	0	0.0%								
No Reported Risk	11	14.5%								
Jurisdictions of HIV Diagnosis v	vith Highe	est Morbidit	у							
Baltimore City	36	47.4%	7.2	Female Male						
Prince George's County	11	14.5%	1.4							
Baltimore County	9	11.8%	1.3	Age at HIV Diagnosis						
Correctional Facility	0	0.0%		IDU HET MSM/IDU MSM						

Note. Correctional facilities are not a high morbidity jurisdiction, but persons residing in correctional facilities are a priority population. Country of birth data not complete for all people.

HIV/HCV Diagnosis: New diagnoses of HIV and HCV within 12 months.

Rate: Number of people living with a HIV/HCV diagnosis divided by the population and multiplied by 100,000.

HCV: Hepatitis C Virus MSM: Male-to-male Sexual Contact IDU: Injection Drug Use HET: Heterosexual Contact					
Center for HIV Surveillance, Epidemiology and Evaluation	 Data source: Maryland Enhanced HIV/AIDS Reporting System & the				
Prevention and Health Promotion Administration	Maryland National Electronic Disease Surveillance System. Data for 2020 should be interpreted with caution due to the impact of				
Maryland Department of Health	the COVID-19 pandemic on access to HIV testing and care-related				
http://phpa.health.maryland.gov/OIDEOR/CHSE	services.				

People Living with HIV Ever Diagnosed with HCV	No.	% of Total	Rate	% with HIV Viral Load Suppressed	HCV Elim Reduce the
Total	3,520	100.0%	69.0	72.1%	Reduce the
Age on December 31, 2020	5,520	100.070	05.0	72.170	201
13-24	4	0.1%	0.4	75.0%	201
25-34	121	3.4%	14.6	49.6%	
35-44	253	7.2%	31.9	53.4%	Reduce the
45-54	665	18.9%	85.6	67.5%	
55+	2,477	70.4%	137.1	76.3%	
Sex at Birth					Among tho
Male	2,352	66.8%	96.2	72.1%	coinfected
Female	1,168	33.2%	44.0	72.1%	and HCV, w
Race/Ethnicity					HIV was the cause of de
Hispanic	110	3.1%	22.5	69.1%	22% had a
Non-Hispanic Black	2,712	77.0%	177.0	73.4%	secondary of
Non-Hispanic White	439	12.5%	16.8	66.1%	death of HC
Non-Hispanic Other	259	7.4%	56.1	69.5%	deathorne
Reported HIV Exposure Cate	egory				
MSM	477	13.6%		72.5%	
IDU	2,046	58.1%		70.9%	HIV Conti
MSM/IDU	284	8.1%		76.4%	
HET	484	13.8%		76.9%	
Perinatal Transmission	7	0.2%		71.4%	
Other	11	0.3%		81.8%	100%
No Reported Risk	211	6.0%		64.9%	80% ب
Jurisdictions of HIV Diagnos	is with Hig	ghest Morbi	dity		60% 60%
Baltimore City	2,110	59.9%	423.0	71.8%	a 40%
Prince George's County	348	9.9%	45.7	72.1%	20%
Baltimore County	423	12.0%	60.6	72.8%	0%
Corrections	34	1.0%		70.6%	0,0

At the end of 2020, there were 3,520 people living with diagnosed HIV who had ever been diagnosed with HCV.

Note. Age-adjusted death rate calculated using 2000 Standard Million Population for the United States. 2019 reported deaths are preliminary.

For additional information on the HCV elimination indicators please see:

https://www.cdc.gov/hepatitis/policy/PDFs/NationalProgressReport.pdf

Reduce the rat	e of repo	rted acut	te HC	V infecti	ons					
2010	th	Among people living with diagnosed HIV, there has been three newly reported acute HCV infection since 2010. Two people were diagnosed in 2010 and 1 in 2019.								
Reduce the rat	e of HCV-	related	death	S						
Among those coinfected with	ו HIV	HCV-Related Age-Adjusted Death Rate Among People Diagnosed with HIV								
and HCV, wher	e te	2016						0.47		
HIV was the leading cause of death,		2017					-	.42		
22% had a	e freed for a second	2018 2019			0.3	21	0.36			
secondary cause of			.00	0.10	0.20	0.30	0.40	0.50		
death of HCV.						th Data	per 100,0	000		
HIV Continu	um of C	are	-7E	sc-Aujus		in nate	501 100,0			
	Diagnos						of Care			
	100.09		ng Pe	ople Ag	ed 13+,	2020				
100%				83.4%	6		72.1%			
۳۵% – ۲ ۵۵۰۲							72.1/0			
60% —										
a 40%										
0%	3,520			2,934	•		2,537			
					_	-				
	HIV Diagn	osed	R	etained i	n Care	Suppre	ssed Viral	Load		

Among current Maryland residents living with diagnosed HIV and ever diagnosed with HCV